

Is It Still Prohibited to Perform Spinal Anesthesia in Patients with Advanced Heart Disease?

Maryam Vosoughian¹, Shideh Dabir¹, Faramarz Mosaffa^{2*}

¹Anesthesiology Research Center, Department of Anesthesiology & Critical Care, Taleghani Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

²Anesthesiology Research Center, Department of Anesthesiology & Critical Care, Akhtar Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

ARTICLE INFO

Article history:

Received 23 May 2024

Revised 15 June 2024

Accepted 28 June 2024

Spinal anesthesia is generally prohibited in patients with severe valvular heart disease (e.g. severe aortic stenosis), cardiomyopathies and heart failure concerning the hemodynamic effects of sympathetic system blockade in patients with limited cardiac output [1-2]. However, based on our experience, in these patients, intrathecal single injection of low doses of local anesthetic (LA) (bupivacaine 5 mg) in combination with 25-50 micrograms of fentanyl or 5 micrograms of sufentanil can be administered safely in lower abdominal and lower limb surgeries lasting 60 to 90 minutes [3-4]. In longer surgeries, we place epidural catheters in the intrathecal space through Tuohy epidural needle and intermittently administer low doses of LA plus low doses of fentanyl or sufentanil as surgically necessary. In these settings, the hemodynamics are stable and the appropriate time covers with no significant adverse effects. We did not observe post-spinal headache in our cases. May be because these patients are mostly in the old age and not at the risk of post-spinal headache.

Standard continuous intrathecal catheters are not available in many centers. So, we suggest that in patients with limited cardiac output, placement of epidural catheters in the spinal space through Tuohy epidural needle can be used as continuous spinal anesthesia for injection of low doses of intrathecal drugs. However, the

risk- benefit ratio should be considered when using spinal anesthesia in this group of patients.

In general, addition of low dose short-acting opioids to low dose LA provides stable hemodynamics, ensures effective anesthesia and enhance postoperative analgesia.

References

- [1] Li M, Huang H. Anesthetic Management of Patients with Dilated Cardiomyopathy Undergoing Noncardiac Surgery. *Medicina*. 2023; 59(9):1567.
- [2] Lopes L, Marialva J, Cardoso H. Spinal Anesthesia in Hemodynamic Instability: A Case Report. *Cureus*. 2023; 15(1):e33821.
- [3] Vosoughian M, Azimaraghi O, Moshari MR, Dahi M, Ghasemi M, Khazaei N. The Effect of Different Combination Doses of Intrathecal Hyperbaric Bupivacaine 0.5% and Sufentanil on the Hemodynamic Profile of Geriatric Patients Undergoing Hip Surgery under Spinal Anesthesia. *Acad J Surg*. 2016; 3(3-4): 42-5.
- [4] Almeida CR, Vieira LS, Cunha P, Gomes A. Low-dose spinal block combined with epidural volume extension in a high-risk cardiac patient: A case-based systematic literature review. *Saudi J Anaesth*. 2022; 16(4):383-389.

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: faramarzmosaffa@sbmu.ac.ir