



Global Prevalence Post-Sternotomy Pain Syndrome (PSPS) after Cardiac Surgery: Systematic Reviews and Meta-Analyse

Mohammad Ali Roozegar^{1,2}, Ali Erfani³, Aminollah Vasigh^{4*}

¹Oral and Dental Health Research Center, Ilam University of Medical Sciences, Ilam, Iran.

²School of Dentistry, Ilam University of Medical Sciences, Ilam, Iran.

³Department of Neurosurgery, School of Medicine Imam Hossein Hospital Shahid Beheshti University of Medical Sciences, Tehran, Iran.

⁴Department of Anesthesiology, Medicine Faculty, Ilam University of Medical Science, Ilam, Iran.

ARTICLE INFO

Article history:

Received 15 August 2024

Revised 03 September 2024

Accepted 17 September 2024

Keywords:

Post-sternotomy pain syndrome;
Cardiac surgery;
Pain;
Systematic reviews and meta-analyses

ABSTRACT

Background: One of the important types of pain is post sternotomy pain syndrome (PSPS). PSPS is defined as pain that persists for 2 months after sternotomy-open cardiac surgery without a specific cause.

Methods: In this meta-analysis study, patients undergoing cardiac surgery in which the prevalence of PSPS has been reported were included in the study. Search keywords included pain, heart surgery, prevalence, heart problems, sternotomy and post-sternotomy pain syndrome. The search was conducted by two researchers based on PICO criteria. The time range of the surge was from the beginning of the publication of articles on the outbreak of PSPS until the beginning of August 2024. Data extraction was done using the researcher's checklist and CMA software was used for data analysis.

Results: In the search that was done with the primary keywords, 108 articles were extracted, after checking the inclusion and exclusion criteria, 8 articles with a sample size of 3423 patients were included in the meta-analysis stage. The analyzed articles were between the years 2001 and 2024, and their methodology was done in different ways, including retrospective, interview, case-control and original. The age range of patients varied from 13 months to 63 years and the sample size of patients varied from 48 patients to 1395 patients. the overall prevalence of PSPS in the 8 analyzed patient articles was 33.4% (with 95% confidence interval: 22.7-46.1).

Conclusions: The prevalence of PSPS in patients worldwide was reported, and drug and non-drug measures are suggested to reduce pain after surgery.

Introduction

Cardiovascular diseases are a common disease in the world, which in addition to physical symptoms such as chest pain, shortness of breath, limitation in activities, also causes mental problems in patients. In fact, heart problems are one of the chronic and high-prevalence diseases that, due to the advances made in the medical field, have led to the hospitalization of a

large number of patients [1-3]. Factors affecting cardiovascular diseases include aging, high blood pressure, body mass index, suffering from other chronic diseases such as diabetes, history of smoking, and hypercholesterolemia [4-5].

In order to improve and reduce the complications of cardiac patients, various methods will be used, including angiography, pacemaker implantation, angioplasty for coronary and cardiac surgeries [6-7]. Annually, in the world, about 2 million people undergo median

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: aminollahvasigh@gmail.com

Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

sternotomy [8]. After heart surgery, the pain is at its most severe in the first 24 hours, and these types of pain decrease over time [9-10].

In all patients, especially patients undergoing heart surgery, pain management is in a special place. Pain after heart surgery may be caused by things including sternal retraction, chest tube or sternotomy [11-12]. One of the important types of pain is post sternotomy pain syndrome (PSPS). PSPS is defined as pain that persists for 2 months after sternotomy-open cardiac surgery without a specific cause. PSPS is an unknown disorder in terms of etiology, which is considered as a common problem [8,12].

PSPS causes the quality of life of patients to decrease due to the pain it causes. If these patients experience pain while performing everyday activities, the pain affects the social, cognitive and psychological aspects of the patient [12-13]. In addition to psychological factors, patients who undergo heart surgery due to emergency conditions are more at risk of chronic pain [14-15]. Factors such as age, gender, anxiety and depression, obesity and history of smoking are effective in aggravating this disease [14-15].

The members of the treatment team, in addition to focusing on the causes and factors affecting the development of the disease, should do their best to meet the mental and physical needs of the patients. The need for pain relief is one of the important needs of patients, and in order to relieve it, the treatment staff should have the necessary information on the prevalence of pain [16-18]. Inadequate pain control of patients after surgery can lead to complications such as lengthening the duration of hospitalization, increasing the rate of disease complications, increasing the rate of death and disability, increasing the costs of hospitalization, the occurrence of disorders in other vital organs of the body, including circulation, blood (increased oxygen consumption and tachycardia), pulmonary (pneumonia and atelectasis) and increased neurohormonal responses as well as muscle weakness in the skeletal muscle system [11-12].

Aim

This meta-analysis study was conducted with the aim of the prevalence of post-sternotomy pain syndrome (PSPS) in patients undergoing cardiac surgery in the world by systematic review and meta-analysis.

Methods

In this meta-analysis study, patients undergoing cardiac surgery in which the prevalence of PSPS has been reported were included in the study. Inclusion criteria included PSPS prevalence reports, all articles published in English worldwide, and availability of full article files. Case report articles and articles whose data were not available were excluded from the study.

Search keywords included pain, heart surgery, prevalence, heart problems, sternotomy and post-sternotomy pain syndrome. Search was done in Google Scholar, SCOPUS, EBSCO, Science Direct, PUBMED and WOS.

The search was conducted by two researchers based on PICO criteria. The time range of the surge was from the beginning of the publication of articles on the outbreak of PSPS until the beginning of August 2024. Data extraction was done using the researcher's checklist (Table 1) and CMA software was used for data analysis.

Results

In the search that was done with the primary keywords, 108 articles were extracted, after checking the inclusion and exclusion criteria, 8 articles with a sample size of 3423 patients were included in the meta-analysis stage. The analyzed articles were between the years 2001 and 2024, and their methodology was done in different ways, including retrospective, interview, case-control and original. The age range of patients varied from 13 months to 63 years and the sample size of patients varied from 48 patients to 1395 patients (Table 1).

The overall prevalence of PSPS in the 8 analyzed patient articles was 33.4% (with 95% confidence interval: 22.7-46.1) (Figure 2).

Table 1- Specifications of the articles

-	Author	Years	Country	Sampling method	Age, M(SD)	N	%
1	El Nasr et al [19]	2017	Egypt	Retrospective	39.77(9.44) Y	48	12.5%
2	Sudin et al [20]	2024	Malaysia	Retrospective	56(11) Y	1395	20.35%
3	Meyerson et al[21]	2001	Sweden	-	13(11) M	318	28%
4	Lahtinen et al [22]	2006	Finland	Prospective follow-up	58(7) Y	213	48%
5	Kalso et al [23]	2001	Finland	Interviewed	-	720	28%
6	Kalliomäki et al [24]	2016	Finland	-	68.7 (8.0) Y	90	15.4%
7	Rashidi et al [25]	2012	Netherlands	-	63.2(12.2) Y	186	83%

8 Costa et al [26] 2015 Brazil Case-control CPSP patients: 453 39.29%
 58(10.9) Y Control patients:62(12.6) Y




Figure1- Prevalence of PSPS




Figure 2- Funnel Plot of PSPS

Commented [A1]: در متن مشخص شود

Discussion

The prevalence of pain in chronic patients is high, which leads to disability in these patients [27]. According to the results of the studies, it is necessary to pay attention to the etiology, prevalence, effective factors and ways to reduce pain in patients [28-32, 42-45]. The purpose of this study was the prevalence of PSPS by meta-analysis in patients of all countries of the world.

The overall prevalence of PSPS in 8 analyzed articles with a sample size of 3423 patients was 33.4%. The prevalence of pain in different patients has been done in different methodological ways. In this study, the prevalence of pain in different parts of the body and

different diseases will be compared with the results of this study.

Comparison of the results with the group under 18 years old:

In one of the studies reviewed in this meta-analysis study (study by Meyerson et al [21]), the prevalence of PSPS in the age range of 13 months was equal to 28%. While in the meta-analysis study by Karimiyarandi et al. in children and adolescents, the rate of pain in wrists/hands was reported as 6.6%, pain in shoulders as 26.9%, pain in elbows as 2.9%, and pain in knees as 6.6% [31]. In the meta-analysis study by Mandrekar et al., with a sample size of 14,743 people, the prevalence of musculoskeletal chest pain was 16%, and the prevalence in the European continent was 17% [33]. In the study of

Sposito et al., which investigated the prevalence of pain in infants, 50.3% of infants experienced pain and 32.5% of them used drug and non-drug programs for pain relief [34]. It seems that pain is experienced in different patients in this age range, but its prevalence may have differences.

Comparison of the results with the group over 18 years old:

In some of the studies included in the meta-analysis study (Rashidi et al [25] and Kalliomäki et al [24]) the patients were in pre-elderly age or in old age who had experienced pain. In the meta-analysis study of Fayaz et al., the prevalence of chronic pain increased with age. So that in the age range of 18 to 22 years, the prevalence of pain was reported to be about 14.3% and for people over 75 years old, about 62% [35]. Also, in French et al.'s study, the prevalence of neuropathic pain in patients with osteoarthritis was reported as 23% by meta-analysis [36], in the meta-analysis study by Sá et al., there were 12 articles with a sample size of 29,902 patients, the prevalence of chronic pain was 18% [37], In the meta-analysis study by Mansfield et al., it was reported that it varied from zero to 24% with a total rate of 10.6% [38].

Comparison of results with patients reported in the field of pain syndromes:

The overall prevalence of PSPS was 33.4%. In the study of Ezzati et al., the prevalence of Myofascial Pain Syndrome was investigated and it was shown that it was 88.9% and was related to the severity of disability [39]. In Berry et al.'s study, the prevalence of Symptoms of Bladder Pain was reported as 6.53% [40], in Alvarez-Nemegyei et al.'s study, the prevalence of rheumatic pain in 12,686 patients was reported as 5% [41]. The results of this study are consistent with the results of pain prevalence.

Conclusions

The prevalence of PSPS in patients worldwide was reported, and drug and non-drug measures are suggested to reduce pain after surgery.

References

- [1] Kazeminezhad B, Tarjoman A, Borji M. Relationship Between Praying and Self-Care in Elderly with Heart Failure: A Cross-Sectional Study in West of Iran. *J Relig Health*. 2020; 59(1):19-28.
- [2] Borji M, Mousavimoghadam SR, Salimi E, Otaghi M, Azizi Y. The Impact of Spiritual Care Education on Anxiety in Family Caregivers of Patients with Heart Failure. *J Relig Health*. 2019; 58(6):1961-9.
- [3] Khoshkesht S, Ghiyasvandian S, Esmaili M, Rahimzadeh M, Rahmanian M, Zakerimoghadam M. The Effect of Designed Transitional Care Program on Anxiety and General Comfort of the Patients Undergoing Coronary Artery Bypass Graft during Transfer from Open Heart Surgery Intensive Care Unit to General Ward: A Clinical Trial. *Scientific JKUMS*. 2023; 28(5):108-21.
- [4] Janjani P, Salehabadi Y, Motevaseli S, Heidari Moghadam R, Siabani S, Salehi N. Comparison of Risk Factors, Prevalence, Type of Treatment, and Mortality Rate for Myocardial Infarction in Diabetic and Non-diabetic Older Adults: A Cohort Study. *Iranian J Ageing*. 2023; 18(2):268-83.
- [5] Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. *J Am Coll Cardiol*. 2020; 76(25):2982-3021.
- [6] Shin D, Singh M, Shlofmitz E, Scheller B, Latib A, Kandzari DE, et al. Paclitaxel-coated versus sirolimus-coated balloon angioplasty for coronary artery disease: A systematic review and meta-analysis. *Catheter Cardiovasc Interv*. 2024.
- [7] Bradić J, Andjić M, Novaković J, Jeremić N, Jakovljević V. Cardioplegia in Open Heart Surgery: Age Matters. *J Clin Med*. 2023; 12(4).
- [8] Costa MA, Trentini CA, Schafranski MD, Pipino O, Gomes RZ, Reis ES. Factors Associated With the Development of Chronic Post-Sternotomy Pain: a Case-Control Study. *Braz J Cardiovasc Surg*. 2015; 30(5):552-6.
- [9] Gerbershagen HJ, Aduckathil S, van Wijck AJ, Peelen LM, Kalkman CJ, Meissner W. Pain intensity on the first day after surgery: a prospective cohort study comparing 179 surgical procedures. *Anesthesiology*. 2013; 118(4):934-44.
- [10] Bjørnnes AK, Rustoen T, Lie I, Watt-Watson J, Leegaard M. Pain characteristics and analgesic intake before and following cardiac surgery. *Eur J Cardiovasc Nurs*. 2016; 15(1):47-54.
- [11] Cogan J. Pain management after cardiac surgery. *Semin Cardiothorac Vasc Anesth*. 2010; 14(3):201-4.
- [12] Nagaraja PS, Ragavendran S, Singh NG, Asai O, Bhavya G, Manjunath N, Rajesh K. Comparison of continuous thoracic epidural analgesia with bilateral erector spinae plane block for perioperative pain management in cardiac surgery. *Ann Card Anaesth*. 2018; 21(3):323-7.
- [13] Sudin A, Chong C, Hassan R. Incidence and Factors Associated With Post-Sternotomy Pain Syndrome in the National Heart Institute, Malaysia. *J Cardiothorac Vasc Anesth*. 2024; 38(2):466-74.
- [14] van Gulik L, Janssen LI, Ahlers SJ, Bruins P, Driessen AH, van Boven WJ, et al. Risk factors for chronic thoracic pain after cardiac surgery via sternotomy. *Eur J Cardiothorac Surg*. 2011; 40(6):1309-13.
- [15] Bordoni B, Marelli F, Morabito B, Sacconi B, Severino P. Post-sternotomy pain syndrome following cardiac surgery: case report. *J Pain Res*. 2017; 1163-9.

- [16] Mohammadi HR, Erfani A, Jamshidbeigi Y, Rahmatian A, Otaghi M. Effect of using rituximab on disability in patients with multiple sclerosis. *J Med Pharm Chem Res.* 2024; 6(12):1854-60.
- [17] Hatefi M, Komlakh K. Investigating the effect of methylprednisolone pulse on treatment of back pain. *Eur Chem Commun.* 2022; 4(5):366-371.
- [18] Shaker MR, Momennasab M, Dehghanrad F, Dokoohaki R, Dakhesh R, Jaber A. The effect of using augmentative and alternative communication strategies on the quality of nurses' communication with patients with endotracheal intubation in the cardiac surgery intensive care unit. *J Hayat.* 2022; 28(1):102-16.
- [19] El Nasr MMA, Taha A. Persistent post sternotomy chest pain: Does sternal wire removal have a role? *J Egypt Soc Cardio-Thorac Surg.* 2017; 25(2):142-6.
- [20] Sudin A, Chong C, Hassan R. Incidence and factors associated with post-sternotomy pain syndrome in the National Heart Institute, Malaysia. *J Cardiothorac Vasc Anesth.* 2024; 38(2):466-74.
- [21] Meyerson J, Thelin S, Gordh T, Karlsten R. The incidence of chronic post-sternotomy pain after cardiac surgery—a prospective study. *Acta Anaesthesiol Scand.* 2001; 45(8):940-4.
- [22] Lahtinen P, Kokki H, Hynynen M. Pain after cardiac surgery: a prospective cohort study of 1-year incidence and intensity. *Anesthesiology.* 2006; 105(4):794-800.
- [23] Kalso E, Mennander S, Tasmuth T, Nilsson E. Chronic post-sternotomy pain. *Acta Anaesthesiol Scand.* 2001; 45(8):935-9.
- [24] Kalliomäki ML, Puolakka P, Huhtala H, Sisto T, Järvelä K. Bedside diagnosis of persistent post-sternotomy pain. *Acta Anaesthesiol Scand.* 2016; 60(7):969-76.
- [25] Rashidi S, Elenbaas TW, Hamad MAS, van Suijlekom HJ, van Straten AH. Does removal of steel wires relieve post-sternotomy pain after cardiac surgery?. *Asian Cardiovasc Thorac Ann.* 2013; 21(4):409-13.
- [26] Costa MACd, Trentini CA, Schafranski MD, Pipino O, Gomes RZ, Reis ESdS. Factors associated with the development of chronic post-sternotomy pain: a case-control study. *Rev Bras Cir Cardiovasc.* 2015; 30(5):552-6.
- [27] Erfani A, Aghamiri SH, Karimi R. Investigating the Role of Hyper Density Signal Length in the Middle Cerebral Artery on the Degree of Disability of Arterial Ischemic Stroke Patients. *Arch Neurosci.* 2024; 11(4):e150888.
- [28] Rahmatian A, Bastani E, Shokri F, Karbasfrushan A. Prevalence of Hemiplegic Shoulder Pain in Iran: A Systematic Review and Meta-analysis. *Anesth Pain Med.* 2023; 13(3):e136423.
- [29] Bastani E, Rizehbandi M, Shokri F. Prevalence of Pain and Factors Affecting it in Patients with Lung Cancer in Ilam. *Int J Cancer Manag.* 2024; 17(1):e136637.
- [30] Hatefi M, KomLakh K. Investigation of the effect of Duloxetine on pain status of patients with spinal cord injuries: A systematic review of drug therapy. *Eur Chem Commun.* 2022; 4(3):256-62.
- [31] Karimiyarandi H, Karbasfrushan A. Effect of duloxetine on pain relief in patients undergoing arthroplasty surgery: a systematic review. *Gomal J Med Sci.* 2022; 20(4):213-20.
- [32] Mohammadi HR, Asadoola Y, Erfani A, Amin NG, Karimiyarandi H, Sadeghi S, Abiri M. Effectiveness of Pulse Intravenous Infusion of Methylprednisolone on Pain in Patients with Lumbar Disc Herniation: A Randomized Controlled Trial. *Anesth Pain Med.* 2024; 14(4): e149442.
- [33] Mandrekar S, Venkatesan P, Nagaraja R. Prevalence of musculoskeletal chest pain in the emergency department: a systematic review and meta-analysis. *Scand J Pain.* 2021; 21(3):434-44.
- [34] Sposito NPB, Rossato LM, Bueno M, Kimura AF, Costa T, Guedes DMB. Assessment and management of pain in newborns hospitalized in a Neonatal Intensive Care Unit: a cross-sectional study. *Rev Latino-Am Enfermagem.* 2017; 25:e2931.
- [35] Fayaz A, Croft P, Langford R, Donaldson L, Jones G. Prevalence of chronic pain in the UK: a systematic review and meta-analysis of population studies. *BMJ open.* 2016; 6(6):e010364.
- [36] French HP, Smart KM, Doyle F, editors. Prevalence of neuropathic pain in knee or hip osteoarthritis: a systematic review and meta-analysis. *Semin Arthritis Rheum.* 2017; 47(1): 1-8.
- [37] Sá KN, Moreira L, Baptista AF, Yeng LT, Teixeira MJ, Galhardoni R, et al. Prevalence of chronic pain in developing countries: systematic review and meta-analysis. *Pain reports.* 2019; 4(6):e779.
- [38] Mansfield KE, Sim J, Jordan JL, Jordan KP. A systematic review and meta-analysis of the prevalence of chronic widespread pain in the general population. *Pain.* 2016; 157(1):55-64.
- [39] Ezzati K, Ravarian B, Saberi A, Salari A, Reyhanian Z, Khakpour M. Prevalence of cervical myofascial pain syndrome and its correlation with the severity of pain and disability in patients with chronic non-specific neck pain. *Arch Bone Joint Surg.* 2021; 9(2):230.
- [40] Berry SH, Elliott MN, Suttorp M, Bogart LM, Stoto MA, Eggers P, et al. Prevalence of symptoms of bladder pain syndrome/interstitial cystitis among adult females in the United States. *J Urol.* 2011; 186(2):540-4.
- [41] Alvarez-Nemegyei J, Pelaez-Ballestas I, Rodriguez-Amado J, Sanin LH, Garcia-Garcia C, Garza-Elizondo MA, et al. Prevalence of rheumatic regional pain syndromes in adults from Mexico: a community survey using COPCORD for screening and syndrome-specific diagnostic criteria. *J Rheumatol Suppl.* 2011; 86:15-20.
- [42] Kloping LP, Utomo DN, Setiawati R, Suryawati S, Irianto KA. Knee spine syndrome. paradigm to

- clinical decision upon patient's pain perception. *Journal of Medicinal and Pharmaceutical Chemistry Research*. 2025; 7(6):1086-95.
- [43] Vellaiyan A. Comparative effects of hegu point ice massage and lidocaine gel on arteriovenous fistula puncture related to pain in hemodialysis patients: A randomized control trail. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 2025; 7(4), 618626.
- [44] Ghiasi N, Reza Mohammadi H, Karimiyarandi H, Otaghi M. Associations between physical activity and pain with self-efficacy in the elderly with chronic pain. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 2025; 7(1): 17.
- [45] Wardana A, Sumartono Ch, Sulisty Putri H, Maskoep WI. The effectiveness of opioid treatment in patients with chronic cancer pain experiencing breakthrough cancer pain (BTcP): Biomarker evaluation in the tetrahydrobiopterin pathway. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 20, 6(11), 16431654.