

Risk Factors Affecting Deep Vein Thrombosis in Pregnant Women

Shideh Ariana¹, Mohammad Karimian², Behzad Azimi^{3*}

¹Department of Obstetrics and Gynecology, School of Medicine, Preventative Gynecology Research Center, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

²Department of Surgery, School of Medicine Emam Khomeini Hospital, Ilam University of Medical Sciences, Ilam, Iran.

³Department of General Surgery, School of Medicine, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

ARTICLE INFO

Article history:

Received 07 February 2025

Revised 28 February 2025

Accepted 11 March 2025

Keywords:

Deep vein thrombosis;

Risk factors;

Pregnant women

ABSTRACT

Background: One of the groups at risk for Deep Vein Thrombosis (DVT) is pregnant women. Considering the importance of pregnancy complications, the aim of this study was to determine the risk factors affecting DVT in the ICU.

Methods: In this study, pregnant women admitted to the ICU were included in the study, and instruments used in this study included a registry checklist that included questions. In this study, the clinical records of people with DVT were placed in 2 groups. The sample size included 28 patients with a diagnosis of DVT (case group) and 33 patients without a diagnosis of DVT (control group). Then, the data were analysed with SPSS 16.

Results: Results showed, in the 28 patients studied for whom the diagnosis of DVT was confirmed, the M(SD) of age status was 37.64 (2.57). There was also a significant association between DVT status with a history of thrombocytopenia, chronic diseases, diabetes, and high blood pressure (BP). ($P < 0.05$).

Conclusion: In DVT, variables such as chronic diseases, diabetes, high blood pressure, and thrombocytopenia were effective. For this reason, it is essential to pay due attention to the factors affecting the development of DVT in clinical care for these patients.

Introduction

Deep Vein Thrombosis (DVT) is the occurrence of an unusual blood clot in a vein of the lower limb, which can be a serious threat to a person's life due to its transformation into a pulmonary embolism, which is called pulmonary thromboembolism. Thromboembolism includes DVT and pulmonary embolism. Factors such as venous stasis, increased coagulability, and vascular vulnerability are effective in its development [1-2]. DVT is the occurrence of an unusual blood clot in the lower limb, which if this clot enters the lung, poses a serious threat to the person [1-2].

Risk factors for DVT in trauma centers include spinal cord injury, limb and pelvic fractures, advanced age, blood transfusions, surgery and severe head trauma, previous history of DVT, venous catheterizations, malignancies, immobilization, surgical incision size, hip replacement, shoulder replacement, use of oral contraceptives, infectious diseases, respiratory failure, obesity, varicose veins, pregnancy, obesity and hormones, some of which are modifiable or preventable [2-5].

The goal of treating DVT in the acute phase includes reducing or eliminating pain and swelling of the affected limb, preventing clot spread and pulmonary embolism, and preventing recurrent thrombosis [6]. Given that the

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: azimibehzad1365@gmail.com

DOI:

Copyright © 2025 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

symptoms of DVT are nonspecific, other tests are recommended to confirm or rule out it. In patients suspected of DVT, performing a D-Dimer test is decisive. Also, if the D-Dimer test results are abnormal, color Doppler ultrasound is necessary [7-8].

One of the groups at risk for DVT is pregnant women. The complications of DVT during pregnancy include ulceration, edema, recurrent thrombosis, and skin changes [9-12]. During pregnancy, due attention should be paid to the health of mothers. In fact, various complications occur during pregnancy for the mother, which is why due attention should be paid to their health [13-16].

Aim

Considering the importance of pregnancy complications, the aim of this study was to determine the risk factors affecting DVT in the ICU.

Methods

This study is part of Iran ICU Registry (IICUR) is a Persian ICU-based registry which was launched in 2018 through a collaboration with The Australian and New Zealand ICU (ANZICS). IICUR was approved by Ethics Committee of Shiraz University of Medical Sciences (Ethic Number IR.SUMS.REC.1397.559), and recognized by Iran Ministry of Health as the first and single registry of adult ICU in Iran.

In this study, pregnant women admitted to the ICU were included in the study. Inclusion criteria included 1- gestational age of more than 3 months or admission to the ICU after delivery, 2- hospitalization for more than 48 hours in the ICU, 3- vascular consultation for the

hospitalized patients, and 3- consent of the patients to participate in the study. Exclusion criteria included the presence of motor problems and lower limb defects before admission.

The instruments used in this study included a registry checklist that included questions on age, smoking (yes/no), history of chronic diseases, diabetes (yes/no), hypertension (yes/no), thrombocytopenia (yes/no), amniotic infection (yes/no), thrombophilia (yes/no), acid-base disorders (yes/no), heart disease (yes/no), sickle cell anemia (yes/no), multiple pregnancy (yes/no), antepartum hemorrhage (yes/no), history of surgery in the last 3 months (yes/no), preeclampsia (yes/no), prolonged pregnancy (yes/no), macrosomia (yes/no), fetal growth restriction (yes/no), increased amniotic fluid (yes/no), extensive postpartum infections (yes/no), and cesarean section (yes/no), and results of INR and PTT tests. The instrument questions were derived from previous studies [17-18].

In this study, the clinical records of people with DVT were placed 2 group. The sample size included 28 patients with a diagnosis of DVT (Case group) and 33 patients without a diagnosis of DVT (Control group). Then, the data were analysis with SPSS 16.

Results

Result showed, in the 28 patients studied for whom the diagnosis of DVT was confirmed, the M(SD) of age status was 37.64 (2.57). There was also a significant association between DVT status with History of Thrombocytopenia, chronic diseases Diabetes and High blood pressure (BP). ($P<0.05$) (Table1).

Table1- Factors affecting the occurrence of risk factors associated with DVT(N=28)

Variable	Yes	No
Smoking	0(0)	28(100)
History of chronic diseases Diabetes	11(39.3)	17(60.7)
High blood pressure	7(25)	21(75)
Thrombocytopenia	5(17.9)	23(82.1)
Amniotic infection	1(3.6)	27(96.4)
Thrombophilia	3(10.7)	25(89.3)
Acid-base disorders	1(3.6)	27(96.4)
Heart disease	1(3.6)	27(96.4)
Sickle cell anemia	1(3.6)	27(96.4)
Multiple pregnancies	3(10.7)	25(89.3)
Prepartum hemorrhage	4(14.3)	24(85.7)
History of surgery in the last 3 months	3(10.7)	25(89.3)
Preeclampsia	1(3.6)	27(96.4)
Prolonged pregnancy	0(0)	28(100)
Macrosomia	1(3.6)	27(96.4)
Incomplete fetal growth	0(0)	28(100)
Increased amniotic fluid	0(0)	28(100)
Extensive postpartum infections	1(3.6)	27(96.4)
Cesarean section	0(0)	28(100)
Partial thromboplastin time (PTT), M(SD)	40.1	22.9

International normalized ratio (INR) M(SD)	1.74	0.73
Age	37.64	2.57

Discussion

Attention to the concept of health, whether it is the prevention of diseases or their treatment, is one of the priorities of the health care system. For this reason, conducting epidemiological studies in order to identify the causes of diseases and the factors affecting them is an important necessity [19-24]. On the other hand, attention to the concept of health in pregnant women is one of the most important priorities of the health care system that should be given due attention [25]. According to the findings of this study, various factors are effective in the development of DVT. These factors include Diabetes, High BP and Thrombocytopenia.

In the study by Anvari et al., it was showed that the factors affecting DVT include age over 60 years, body mass index and previous history of DVT [26]. Also, in the study by Zhang et al., it was shown among the women surveyed that in the first trimester of pregnancy, the factors affecting DVT included a previous history of DVT, immune diseases, and thrombophilia [27]. In the study by Mugeni et al., the prevalence of proximal DVT in hospitalized patients was reported to be 5.5%. In this study, 7% of them were pregnant. Also, the factors affecting DVT include a previous history of DVT, recent long-term travel, long-term immobility, and active malignancy [28]. Also, in the study of Mallah et al., it was shown that diabetes, cardiovascular diseases, hypertension, thrombocytopenia and history of surgery in the last 3 months were among the factors affecting the occurrence of DVT [17].

In a study by Mahjobipoor et al., 2000 records of patients admitted to the ICU were retrospectively reviewed, and 20 patients were diagnosed with DVT. According to the findings, DVT in patients was significantly associated with patient age, type of hospitalization (surgical or internal), nutritional status, length of stay, and Apache score ($P < 0.05$) [29]. In a study by Miri et al., 1387 patient records were reviewed, 500 patients were diagnosed with DVT. The mean age of the patients was 60 ± 18 years. In addition to older age, the length of stay in the ICU was also associated with the development of DVT [30].

Other studies include the study by Javadi Moghaddam. In this study, all patients referred to the hospital were examined and it was shown that most patients with DVT were in the age range of 30-40 years [31]. In the study by Mousa et al., in most of the patients studied, factors such as immobility, stroke, fracture, chemotherapy, obesity, oral contraceptive use, and malignancies were found to be important factors in the development of DVT [32]. In the study by Takeda et al., patients hospitalized for neurodegenerative diseases were examined and the rate

of DVT was reported to be higher in patients with female gender and higher blood pressure [33].

Conclusion

In DVT, variables such as chronic diseases Diabetes, High blood pressure and Thrombocytopenia were effective. For this reason, it is essential to pay due attention to the factors affecting the development of DVT in clinical care for these patients.

References

- [1] Kraaijpoel N, van Es N, Porreca E, Büller HR, Di Nisio M. The diagnostic management of upper extremity deep vein thrombosis: A review of the literature. *Thromb Res.* 2017; 156:54-9.
- [2] Di Minno MN, Ambrosino P, Ambrosini F, Tremoli E, Di Minno G, Dentali F. Prevalence of deep vein thrombosis and pulmonary embolism in patients with superficial vein thrombosis: a systematic review and meta-analysis. *J Thromb Haemost.* 2016;14(5):964-72.
- [3] Nicholson M, Chan N, Bhagirath V, Ginsberg J. Prevention of venous thromboembolism in 2020 and beyond. *J Clin Med.* 2020;9(8):2467.
- [4] Geersing G, Zuithoff N, Kearon C, Anderson D, Ten Cate-Hoek A, Elf J, et al. Exclusion of deep vein thrombosis using the Wells rule in clinically important subgroups: individual patient data meta-analysis. *Bmj.* 2014;348.
- [5] Naseri A, Dehghani A, Moharrami MR. Prevalence of deep vein thrombosis following knee replacement surgery in patients with a history of chemotherapy. *Iran J Orthop Surg.* 2021;
- [6] Gram AS, Bladbjerg EM, Skov J, Ploug T, Sjødin A, Rosenkilde M, et al. Three months of strictly controlled daily endurance exercise reduces thrombin generation and fibrinolytic risk markers in younger moderately overweight men. *Eur J Appl Physiol.* 2015;115(6):1331-8.
- [7] Niimi K, Nishida K, Lee C, Ikeda S, Kawai Y, Sugimoto M, et al. Optimal D-Dimer cutoff values for diagnosing deep vein thrombosis in patients with comorbid malignancies. *Ann Vasc Surg.* 2024; 98:293-300.
- [8] Kitamura F, Shiraishi Y, Sakata K, Takata N, Harada K, Yoshinaka I, et al. Screening for Deep Vein Thrombosis Using D-dimer Levels Based on Surgical Patients' Characteristics. *Cureus.* 2024;16(12):e75565.
- [9] Kujovich JL. Hormones and pregnancy: thromboembolic risks for women. *Br J Haematol.* 2004;126(4):443-54.
- [10] James AH, Jamison MG, Brancazio LR, Myers ER.

- Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *Am J Obstet Gynecol.* 2006;194(5):1311-5.
- [11] James AH, Tapson VF, Goldhaber SZ. Thrombosis during pregnancy and the postpartum period. *Am J Obstet Gynecol.* 2005;193(1):216-9.
 - [12] Devis P, Knuttinen MG. Deep venous thrombosis in pregnancy: incidence, pathogenesis and endovascular management. *Cardiovasc Diagn Ther.* 2017;7(Suppl 3):S309-s19.
 - [13] Putrie SW, Wicaksono B, Hadi THS. Unveiling a rare challenge: Giant cervical polyp in pregnancy: A case report and management insights. *J Med Pharm Chem Res.* 2025;7(9):1970-8.
 - [14] Suherman RWP, Akbar MIA. Maternal and perinatal outcomes in obese pregnant women. *J Med Pharm Chem Res.* 2025;7(7):1336-48
 - [15] Mezaal HD, Al-Fartosi KG, El Arbi M. Effect of sleep disturbance on the hormone melatonin and its relationship to some biochemical analyzes among some pregnant women in Thi-Qar governorate. *J Med Pharm Chem Res.* 2025;7(7):1411-9.
 - [16] Nabilah H, Akbar MIA, Joewono HT. The differences of risk factors and outcomes of pregnancy with obesity who had failed and successful induction of labor in 2023. *J Med Pharm Chem Res.* 2025;7(7):1503-12.
 - [17] Mallah F, Ilkhchi NS, Alvandfar D. The risk factors of deep vein thrombosis in pregnancy intensive care unit, Alzahra Hospital in Tabriz, 2019. *The Iranian Journal of Obstetrics, Gynecology and Infertility.* 2021; 24(11): 10-16.
 - [18] Kafeza M, Shalhoub J, Salooja N, Bingham L, Spagou K, Davies AH. A systematic review of clinical prediction scores for deep vein thrombosis. *Phlebology.* 2017;32(8):516-31.
 - [19] Hatefi M, Parvizi R, Borji M, Tarjoman A. Effect of self-management program on pain and disability index in elderly men with osteoarthritis. *Anesth Pain Med.* 2019;9(4).
 - [20] Mohajerzadeh L, Ebrahimian M, Sarafi M, Ebrahimisaraj G, Tabatabaei SR, Noorbakhsh SM, et al. Effects of COVID-19 Pandemic on the Frequency of Complicated Appendicitis in Pediatric Populations. *Arch Pediatr Infect Dis.* 2023;11(1).
 - [21] Esmailizadeh AH, Ebrahimisaraj G, Sarafi M, Rouzrokh M, Mohajerzadeh L, Ghoroubi J, et al. The outcome and complications of modern staged repair surgery in newborns with classic bladder exstrophy in different genders: A retrospective study. *Birth Defects Research.* 2023;115(16):1469-74.
 - [22] Ebrahimisaraj G, Sarafi M, Ebrahimian M, Esmailizadeh AH, Ladan AH, Khoshkhabar M, et al. Short-term Outcomes of Pancreatectomy in Congenital Hyperinsulinism: A Retrospective Multi-center Study. *Iran J Neonatol.* 2023;14(2).
 - [23] Mohammadi HR, Erfani A, Sadeghi S, Komlakh K, Otaghi M, Vasig A. Investigating Factors Affecting Mortality Due to Spinal Cord Trauma in Patients Admitted to the Intensive Care Unit. *Bull Emerg Trauma.* 2024;12(3):136-41.
 - [24] Erfani A, Aghamiri SH, Karimi R. Investigating the Role of Hyper Density Signal Length in the Middle Cerebral Artery on the Degree of Disability of Arterial Ischemic Stroke Patients. *Arch Neurosci.* 2024;11(4):e150888.
 - [25] Azami M, Nasirkandy MP, Mansouri A, Darvishi Z, Rahmati S, Abangah G, et al. Global prevalence of helicobacter pylori infection in pregnant women: a systematic review and meta-analysis study. *Int J Womens Health Reprod Sci.* 2017;5(1):30-6.
 - [26] Anvari HM, Zeinalzadeh M, Ilkhchi NS, Alvandfar D. The factors affecting deep vein thrombosis following laparoscopic hysterectomy and the factors affecting it. *The Iranian Journal of Obstetrics, Gynecology and Infertility.* 2021; 24(11):17-23.
 - [27] Zhang C, Zhang X, Liu G. Deep vein thrombosis in early pregnancy: A retrospective study. *Gynecol Obstet Clin Med.* 2023;3(1):44-50.
 - [28] Mugeni R, Nkusi E, Rutaganda E, Musafiri S, Masaisa F, Lewis KL, et al. Proximal deep vein thrombosis among hospitalised medical and obstetric patients in Rwandan university teaching hospitals: prevalence and associated risk factors: a cross-sectional study. *BMJ open.* 2019; 9(11):e032604.
 - [29] Mahjobipoor H, Rahimi-Varposhti M, Shetabi H, Heidari S. Predictive factors of deep vein thrombosis in patients admitted to intensive care unit. *Tehran Univ Med J.* 2021;78(12):828-34.
 - [30] Miri M, Goharani R, Sistanizad M. Deep Vein Thrombosis among Intensive Care Unit Patients; an Epidemiologic Study. *Emerg (Tehran).* 2017;5(1):e13.
 - [31] Javadi Moghaddam M, Bakhshi Ali Abad H. Seasonal Changes in the Occurrence of Deep Vein Thrombosis (DVT) in Patients Referred to Aliebn-e-Abitaleb Hospital of Rafsanjan City in 2017: A Short Report. *J Rafsanjan Univ Med Sci.* 2020;18(11):1183-90.
 - [32] R Mousa B, E Mahdy N, Mahmoud SF, G Mohamed M. Risk factors for deep venous thrombosis among patients admitted to Vascular Unit. *Egyptian Journal of Health Care.* 2022;13(2):270-86.
 - [33] Takeda T, Koreki A, Kokubun S, Saito Y, Ishikawa A, Iose S, et al. Deep vein thrombosis and its risk factors in neurodegenerative diseases: A markedly higher incidence in Parkinson's disease. *J Neurol Sci.* 2024; 457:122896.