

Association of Cardiopulmonary Bypass and Aortic Cross-Clamp Durations with ICU Length of Stay in Cardiac Surgery Patients: A Retrospective Study at Nasiriya Heart Center

Hussein Ali Hussein¹, Myasar Jasim Mohammed², Qusay Abdulazhara Yaqoob³, Mohammed AbdulZahra Sasaa⁴, Majid Fakhir Alhamaidah^{1*}, Hussein Alkhfaji¹, Ammar Hoom Mahdi⁵, Abbass Hussein Haydar¹

¹Department of Anesthesia, College of Health and Medical Technologies, Alayen Iraqi University, Thi-Qar, Iraq.

²Department of General Surgery, College of Medicine, Diyala University, Diyala, Iraq.

³Department of Surgery, College of Medicine, University of Kufa, Al-Kufa Street, Najaf, Iraq.

⁴Department of Anesthesia, College of Health and Medical Techniques, Al-Mustaqbal University, Hillah, Iraq.

⁵Department of Anesthesia, College of Health and Medical Techniques, University of Bilad Alrafidain, Diyala, Iraq.

ARTICLE INFO

Article history:

Received 25 September 2025

Revised 16 October 2025

Accepted 30 October 2025

Keywords:

Cardiopulmonary bypass;
Aortic cross-clamp;
Cardiac surgery;
Postoperative complications;
Intensive care units

ABSTRACT

Background: Cardiopulmonary bypass (CPB) and aortic cross-clamping are essential components of many cardiac surgical procedures. However, prolonged durations have been linked to adverse outcomes, including delayed recovery and increased intensive care unit (ICU) resource utilization. Understanding their impact is critical for improving perioperative management and optimizing patient outcomes.

We aimed to assess how cardiopulmonary bypass time (CPBT) and aortic cross-clamp time (ACCT) affect the length of ICU stay in patients undergoing elective cardiac surgery.

Methods: Our study was retrospective clinical study which conducted at Nasiriya Heart Center, Iraq, between September 2023 and September 2024. A total of 100 patients (aged 18–65 years) who underwent elective cardiac surgery were included. Demographic data, type of surgery, admission and discharge dates, and intraoperative variables (CPBT and ACCT) were collected. Associations between CPBT, ACCT, and ICU stay were analysed by using suitable statistical approaches.

Results: Of the 100 patients, 47% were male and 53% female, with a mean age of 44 years. The mean CPBT was 112.6 ± 44.2 minutes, and the mean ACCT was 69.0 ± 34.5 minutes. Gender showed no significant effect on ICU stay. In contrast, both prolonged CPBT and ACCT were significantly associated with longer ICU stays ($P = 0.047$ and $P = 0.005$, respectively).

Conclusion: Extended CPBT and ACCT are significant predictors of prolonged ICU stay after cardiac surgery. Strategies to minimize intraoperative times may help reduce postoperative ICU occupancy and improve overall resource allocation in cardiac surgical care.

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: Majid.mutar@alayen.edu.iq

DOI:

Copyright © 2026 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

Introduction

Cardiopulmonary bypass (CPB) represents one of the most significant advancements in modern cardiac surgery. It is a specialized form of extracorporeal circulation in which systemic venous blood is redirected from the heart and lungs and routed through an external circuit, ensuring continuous systemic perfusion and oxygenation during surgery [1]. In most procedures, the heart is isolated from the systemic circulation by aortic cross-clamping (ACC), and a cardioplegic solution is administered. This technique allows surgeons to operate in a motionless and bloodless operative field while preserving vital organ function [2–4].

Despite its advantages, prolonged application of CPB and ACC carries a progressive risk of myocardial ischemia due to interruption of coronary perfusion [5]. To counteract ischemic injury, myocardial protection strategies such as cardioplegia, systemic hypothermia, and cardiac perfusion techniques have been employed. Nevertheless, ACC duration remains a major determinant of postoperative myocardial dysfunction and is closely linked with low cardiac output syndrome (LCOS), often necessitating inotropic support, mechanical assistance, and prolonged intensive care unit (ICU) management [6]. Multiple perioperative factors—including patient demographics, surgical type and complexity, duration of CPB, ACC time, renal dysfunction, and postoperative organ impairment—have been studied for their impact on outcomes [7-8]. Among these, prolonged mechanical ventilation (PMV) exceeding seven days has emerged as a significant determinant of mortality, residual disability, and overall length of stay (LOS) [8-9]. However, most evidence originates from international centers, with limited data from Middle Eastern populations.

In Iraq, and particularly at the Nasiriya Heart Center, there remains a paucity of published studies addressing the relationship between intraoperative variables and ICU outcomes. Considering the potential implications for resource utilization and patient prognosis, it is essential to investigate these associations in local populations. Therefore, the present study aims to evaluate the relationship between CPB duration, ACC time, and postoperative ICU stay in individuals who underwent cardiac surgery at the Nasiriya Heart Center.

Methods

Study Design and Setting

This retrospective, single-center study was conducted at the Nasiriya Heart Center for Cardiac Surgery, Dhi Qar, Iraq. The study included adult patients aged 18–65 years who underwent elective cardiac surgery requiring cardiopulmonary bypass (CPB) over a one-year period, between September 2023 and September 2024.

Study Population

Eligible patients were those admitted postoperatively to the intensive care unit (ICU) following elective cardiac surgery performed with CPB support. Exclusion criteria included patients who underwent emergency cardiac surgery, patients who died within the early postoperative days in the ICU, and patients with incomplete or missing medical records.

A total of 104 patient records were initially screened. Four cases were excluded: two patients who died during the early ICU stay and two patients with chronic comorbidities (hypertension and diabetes mellitus). Ultimately, 100 patient records involved in the ending analysis (Figure 1).




Figure 1- Flow diagram of patient selection for the retrospective cohort study.

Data Collection

Patient data were reviewed and extracted from medical records using a structured data collection form. The following parameters were assessed:

- Demographic data: age and sex.
- Preoperative variables: diagnosis, clinical syndrome, left ventricular (LV) function, and risk category.
- Intraoperative variables: CPB duration (measured from initiation to termination) and aortic cross-clamp (ACC) time.
- Postoperative variables: duration of ICU stay, total length of hospital stay (LOS), and in-hospital mortality.

Admission and discharge dates were documented to precisely determine ICU and hospital LOS.

Patient Grouping

Based on ICU length of stay, patients were stratified into three groups: Group I: ICU stay of 1–3 days, Group II: ICU stay of 4–6 days, and Group III: ICU stay exceeding 6 days. Preliminary analysis of the data indicated that the mean ICU stay was 4 days. For the

purposes of this study, a prolonged holding on in the ICU was defined as more than 4 days after elective CABG surgery.

Ethical Considerations

This study was approved by the Ethical Committee of the Dhi Qar Health Office (ethical code: A8/1/2025).

All procedures adhered to the ethical principles outlined in the Iraqi national research standards and international guidelines for studies involving human participants. Patient confidentiality was strictly maintained through anonymization of data prior to analysis.

Statistical Analysis

For statistical analysis, we used SPSS version 22 (SPSS Inc., Chicago, IL, USA). Percentages and frequencies were depended basically in presenting the categorical data, and continuous variables as medians with interquartile ranges.

The Kolmogorov–Smirnov test was applied to assess the normality of data distribution. As normality assumptions were not met for all variables, non-parametric statistical methods were used. Correlations between study variables were evaluated in two statistical methods, the 1st one was depending the Spearman's rank correlation coefficient, and the 2nd one a comparison between groups were performed with the Mann–Whitney U test.

A two-tailed P value < 0.05 was considered statistically significant.

Results

The study included 100 patients who underwent elective cardiac surgery with cardiopulmonary bypass. Their baseline characteristics are shown in (Table 1).

On average, patients were 44.1 years old (median 47), and just over half (53%) were female."

The mean CPB time was 112.6 ± 45.2 minutes (median 110, range 48–242 minutes), and the mean aortic cross-clamp (ACC) time was 69.0 ± 38.1 minutes (median 65, range 4–208 minutes) (Table 2).

The relationship between gender and ICU length of stay was not significant ($p = 0.089$; 2), indicating that male and female patients had similar ICU durations (Table 3).

Increasing age was significantly associated with longer ICU stay (χ^2 test, $p = 0.001$). All patients aged <21 years stayed 1–3 days, whereas among patients >60 years, only 50% stayed 1–3 days; 25% stayed 4–6 days, and 25% stayed >6 days (Table 4).

Longer CPB time was significantly accompanying with extended ICU stay ($p = 0.047$). All patients with CPB <60 minutes stayed 1–3 days, while patients with CPB 180–240 minutes had only 50% staying 1–3 days, 33.3% staying 4–6 days, and 16.7% staying >6 days (Table 5).

Lengthier ACC time was also significantly linked with lengthy ICU stay ($p = 0.005$). Patients with ACC <60 minutes predominantly stayed 1–3 days (88.1%), whereas those with ACC 180–240 minutes all stayed >4 days (Table 6).

Table 1- Descriptive analysis of the patients' characteristics

Variable	N	Mean \pm SD	Median	n (%)
Age (years)	100	44.1 \pm 15.9	47	
Sex: Male	47	-	-	(47%)
Sex: Female	53	-	-	(53%)

Table 2- Descriptive analysis of the patients' intraoperative variables.

Variable	Mean \pm SD	Median	Range
CPB time (min)	112.6 \pm 45.2	110	48–242
ACC time (min)	69.0 \pm 38.1	65	4–208

Table 3- Gender and ICU stay

Gender	ICU Duration 1–3 Day	ICU Duration 4–6 Day	ICU Duration >6 Day	Total	Chi-square test (P value, df)
Male	30 (63.8%)	12 (25.5%)	5 (10.7%)	47	$p = 0.089$, $df = 2$
Female	28 (52.8%)	17 (32.1%)	8 (15.1%)	53	
Total	58 (58.0%)	29 (29.0%)	13 (13.0%)	100	

Table 4- Age and ICU stay

Age Groups	1–3 Days	4–6 Days	>6 Days	Total	Chi-square test (P value, df)
< 21 years	10 (100%)	0 (0%)	0 (0%)	10 (100%)	(0.001; 6)
21–40 years	27 (96%)	1 (4%)	0 (0%)	28 (100%)	
41–60 years	41 (82%)	8 (16%)	1 (2%)	50 (100%)	
> 60 years	6 (50%)	3 (25%)	3 (25%)	12 (100%)	

Table 5- ICU Duration According to CPB Time

CPB Time	1–3 Days	4–6 Days	>6 Days	Total	Chi-square test (P value, df)
< 60 minutes	13 (100%)	0 (0%)	0 (0%)	13 (100%)	(0.047; 6)
60–119 minutes	49 (89.1%)	5 (9.1%)	1 (1.8%)	55 (100%)	
120–179 minutes	19 (73.1%)	5 (19.2%)	2 (7.7%)	26 (100%)	
180–240 minutes	3 (50%)	2 (33.3%)	1 (16.7%)	6 (100%)	

Table 6- ACC time and ICU stay

ACC Time (minutes)	ICU Duration 1–3 Days	ICU Duration 4–6 Days	ICU Duration > 6 Days	Total (n)	Chi-square test (P value, df)
< 60 minutes	37 (88.1%)	5 (11.9%)	0 (0%)	42 (100%)	p = 0.005, df = 6
60–119 minutes	42 (85.7%)	5 (10.2%)	2 (4.1%)	49 (100%)	
120–179 minutes	5 (71.4%)	1 (14.3%)	1 (14.3%)	7 (100%)	
180–240 minutes	0 (0%)	1 (50%)	1 (50%)	2 (100%)	

Discussion

Cardiopulmonary bypass (CPB) is a medical technique used during certain surgical procedures, especially during open-heart surgery. During cardiac surgery with CPB, the patient's venous blood is pumped through an artificial oxygenator and then returned to the body. While CPB is a life-saving procedure, it can also have several physiological effects on the body [10-11].

Cardiopulmonary bypass (CPB) stimulates a systemic inflammatory response, characterized by consequences such as coagulation disorders, leukocytosis, fever, edema, and multi-organ dysfunction [12-13]. The severity of this response tends to increase with prolonged CPB duration. Furthermore, the extent of CPB-related complications has been shown to correlate not only with bypass duration but also with the patient's preoperative clinical condition [14-15].

In this context, the goal of medical professionals is to manage and minimize this response as much as possible to ensure the patient's safety and well-being. Various strategies and techniques, such as utilizing biocompatible materials in the CPB circuit, minimizing the duration of CPB, and administering anti-inflammatory medications, are employed to alleviate the inflammatory effects which allied with CPB [1-2].

Undeniably, aortic cross-clamping is a common component of open cardiac procedure, particularly during procedures which involving bypass grafting of the coronary arteries (CABG). An aortic clamp is used to temporarily stop blood flow through the aorta, allowing the surgeon to work on the heart without interference from blood flow. While aortic cross-clamping is necessary for the surgical procedure, it can also have implications in inflammation (e.g., ischemia-reperfusion injury, myocardial inflammation, systemic effects on lungs and kidneys) [16-18]. To mitigate these effects, medical professionals take measures mainly by minimizing cross-clamp time to reduce the risk of ischemia-reperfusion injury and inflammation, by

resorting to cardioplegia that protects the heart from ischemic injury during the period of cross-clamping and by closely monitoring patients during and after surgery [6].

In the literature, numerous clinical investigations have consistently demonstrated that the lengths of ACC and CPB utilization during cardiac procedures independently influence the risks of both mortality and morbidity. To our knowledge, there is a paucity of such retrospective studies in Iraq. To this purpose, we surveyed the association between cardiopulmonary bypass duration and aortic cross-clamp time with post-cardiac surgery ICU hospitalization in the Nasiriya Heart Center for Cardiac Surgery [19-21].

Our findings had been revealed that the age, prolonged use of CPB and ACC time were correlated with increased ICU stay ($P=0.001^*$, 0.047^* , and 0.005^* , respectively).

As the contrary, the gender seems to be indifferent ($P=0.089$). Nonetheless, a discrepancy was noted in the literature. Bertrand et al. [22] and Moh'd et al. [6] show no link between age and ICU bed occupancy after cardiac surgery, while Rashid et al. [23] noticed a slight prevalence in men over women. The variance in the study outcomes may be due to different reasons of patients with different perioperative variables and morbidity. In addition, cardiovascular disease has long been viewed as a condition that affects men. Even though the age-specific rates of these conditions are elevated in men if compared to women in most age groups, the lifetime risk of developing them is similar for both men and women [24-25].

In this study, the factor "age" is a good predictor of the ICU bed occupancy, as shown in (Table 4) ($P=0.001$). However, L. Santana-Cabrera et al. concluded that no differences were noticed in ICU stay and its relation to the age, while a difference was found in mortality [26]. Similarly, Gavilanes et al. reported that among survivors, age was not associated with ICU length of stay; however, for individuals who died, increasing age was not correlated with prolonged ICU and overall hospital stay

[27]. Another study demonstrated that the increase in mortality with advancing age among ICU patients is not attributable to greater disease severity. Instead, it suggested that ICU mortality rises with age irrespective of treatment intensity [28].

As expected, we found that CPB and ACC times predict strongly the length of stay in ICU with Pearson's $\rho=0.325^{**}/P=0.047$ and $0.302^{**}/P=0.005$, respectively. These observations are in line with former studies. Nazish Alisher et al. prove this association [29].

Similarly, the findings of Szekely et al. were consistent with ours, as they reported a significant association between CPB duration and postoperative mechanical ventilation time [31]. Furthermore, several other studies have identified CPB duration as a risk factor for prolonged ICU stay in both pediatric and adult populations [23,31-32]. Moreover, in the Jan Bucerius et al. study, risk factors enhance by 1.59 times the length of stay in the ICU when the CPB was greater than 120 minutes; along with their result, in our study, the risk factors increase the length of stay in the ICU when the CPB was greater than 119 min [6]. Discordant results were also reported by other publications [33-34].

The discrepancy in findings is most likely attributable to variations in study methodologies and the inclusion of different sets of confounding perioperative variables compared with our study, such as infection, UTI, vasculitis, pneumonia, diabetes, and other inflammatory conditions that are very reliable risk factors determining the ICU and hospital durations as well as affecting mortality and other morbidity outcomes. Our current survey does not incorporate these factors, which explains the clear and relevant association of CPB or ACC times to ICU bed occupancy. Moreover, the categorization of CPB and ACC times (as 30-minute categories) that we adopted in the same way as in the Mehmood et al. study yielded a greater contribution to the ICU length of stay. This points up the importance of the continuous improvements brought in by CBP and ACC technologies to alleviate their post-surgical consequences.

Before we go any further, we will mention to the main of study's limitations. The most important ones are the retrospective design of the study and the fact that it was only conducted at one center. This means the evidence isn't as strong as it could be and, frankly, we can't prove that one thing actually caused another.

In addition, we were unable to assess certain postoperative outcomes, such as non-infectious pulmonary complications, pneumothorax, cardiomyopathy, shock, mediastinitis, non-union of the sternum wound, pneumonia, infection, lymphedema, duration of mechanical ventilation, and failure of intubation removal. This is attributed to many reasons such as incomplete reports, neither mortality nor survival, as the number of individuals who died was small and not sufficient for any calculation.

Thus, further studies involving a larger set of patients are required to confirm the presented findings. Lastly, as prolonged ICU stay is associated with poor postoperative outcomes and increased financial burden for patients, further Iraqi studies and more efforts are of utmost importance to investigate variables also prone to affect the duration of CBP, aortic clamping, and the ICU discharge after cardiac surgeries.

Conclusion

Our conclusions show that patients who are older or spend more time on cardiopulmonary bypass and aortic cross-clamp tend to elongated stay in the ICU. Given that extended ICU stays are linked to poorer recovery and more complications, it makes sense to factor in ACCT and CPBT when managing patients after surgery.

References

- [1] Stoney WS. Evolution of cardiopulmonary bypass. *Circulation*. 2009; 119(21):2844-53.
- [2] Sarkar M, Prabhu V. Basics of cardiopulmonary bypass. *Indian J Anaesth*. 2017; 61(9):760-767.
- [3] Hessel II EA. What's new in cardiopulmonary bypass. *J Cardiothorac Vasc Anesth*. 2019; 33(8):2296-326.
- [4] Machin D, Allsager C. Principles of cardiopulmonary bypass. *Contin Educ Anaesth Crit Care Pain*. 2006; 6(5):176-81.
- [5] Komagamine M, Fukunishi T, Yamasaki Y, Tomita M, Kinebuchi S, Tomimoto D, et al. Patient Outcomes after Introduction of Novel Myocardial Protection Protocol for Prolonged Aortic Cross-Clamping. *Ann Thorac Cardiovasc Surg*. 2025; 31(1):25-00079.
- [6] Moh'd AF, Al-Odwan HT, Altarabsheh S, Makahleh ZM, Khasawneh MA. Predictors of aortic clamp time duration and intensive care unit length of stay in elective adult cardiac surgery. *Egypt Heart J*. 2021; 73(1):92.
- [7] Ortega-Loubon C, Fernández-Molina M, Pañeda-Delgado L, Jorge-Monjas P, Carrascal Y. Predictors of Postoperative Acute Kidney Injury after Coronary Artery Bypass Graft Surgery. *Braz J Cardiovasc Surg*. 2018; 33(4):323-329.
- [8] De Santo LS, Rubino AS, Montella AP, Golini Petrarcone C, Palmieri L, Galbiati D, et al. Incidence and risk factors of acute kidney injury in redo cardiac surgery: a single center analysis. *Sci Rep*. 2024; 14(1):27267.
- [9] Wang Q, Tao Y, Zhang X, Xu S, Peng Y, Lin L, et al. The Incidence, Risk Factors, and Hospital Mortality of Prolonged Mechanical Ventilation among Cardiac Surgery Patients: A Systematic Review and Meta-Analysis. *Rev Cardiovasc Med*. 2024; 25(11):409.
- [10] Banerjee D, Feng J, Sellke FW. Strategies to

- attenuate maladaptive inflammatory response associated with cardiopulmonary bypass. *Front Surg.* 2024; 11:1224068.
- [11] Rubens FD, Mesana T. The inflammatory response to cardiopulmonary bypass: a therapeutic overview. *Perfusion.* 2004; 19 Suppl 1:S5-12.
- [12] Landis RC, Brown JR, Fitzgerald D, Likosky DS, Shore-Lesserson L, Baker RA, et al. Attenuating the Systemic Inflammatory Response to Adult Cardiopulmonary Bypass: A Critical Review of the Evidence Base. *J Extra Corpor Technol.* 2014; 46(3):197-211.
- [13] Bronicki RA, Hall M. Cardiopulmonary Bypass-Induced Inflammatory Response: Pathophysiology and Treatment. *Pediatr Crit Care Med.* 2016; 17(8 Suppl 1):S272-8.
- [14] Li S, Price R, Phiroz D, Swan K, Crane TA. Systemic inflammatory response during cardiopulmonary bypass and strategies. *J Extra Corpor Technol.* 2005; 37(2):180-8.
- [15] Axtell AL, Fiedler AG, Melnitchouk S, D'Alessandro DA, Villavicencio MA, Jassar AS, et al. Correlation of cardiopulmonary bypass duration with acute renal failure after cardiac surgery. *J Thorac Cardiovasc Surg.* 2020; 159(1):170-178.e2.
- [16] Gelman S. The pathophysiology of aortic cross-clamping and unclamping. *Anesthesiology.* 1995; 82(4):1026-60.
- [17] Zammert M, Gelman S. The pathophysiology of aortic cross-clamping. *Best Pract Res Clin Anaesthesiol.* 2016; 30(3):257-69.
- [18] Yeung KK, Groeneveld M, Lu JJ, van Diemen P, Jongkind V, Wisselink W. Organ protection during aortic cross-clamping. *Best Pract Res Clin Anaesthesiol.* 2016; 30(3):305-15.
- [19] Doenst T, Berretta P, Bonaros N, Savini C, Pitsis A, Wilbring M, et al. Aortic cross-clamp time correlates with mortality in the mini-mitral international registry. *Eur J Cardiothorac Surg.* 2023; 63(6):ezad147.
- [20] Swinkels BM, Ten Berg JM, Kelder JC, Vermeulen FE, Van Boven WJ, de Mol BA. Effect of aortic cross-clamp time on late survival after isolated aortic valve replacement. *Interact Cardiovasc Thorac Surg.* 2021; 32(2):222-228.
- [21] Hu J, Liu Y, Huang L, Song M, Zhu G. Association between cardiopulmonary bypass time and mortality among patients with acute respiratory distress syndrome after cardiac surgery. *BMC Cardiovasc Disord.* 2023; 23(1):622.
- [22] Delannoy B, Guye ML, Slaiman DH, Lehot JJ, Cannesson M. Effect of cardiopulmonary bypass on activated partial thromboplastin time waveform analysis, serum procalcitonin and C-reactive protein concentrations. *Crit Care.* 2009; 13(6):R180.
- [23] Nadeem R, Agarwal S, Jawed S, Yasser A, Altahmody K. Impact of Cardiopulmonary Bypass Time on Postoperative Duration of Mechanical Ventilation in Patients Undergoing Cardiovascular Surgeries: A Systemic Review and Regression of Metadata. *Cureus.* 2019; 11(11):e6088.
- [24] Albrektsen G, Heuch I, Løchen ML, Thelle DS, Wilsgaard T, Njølstad I, et al. Lifelong Gender Gap in Risk of Incident Myocardial Infarction: The Tromsø Study. *JAMA Intern Med.* 2016; 176(11):1673-1679.
- [25] Leening MJ, Ferket BS, Steyerberg EW, Kavousi M, Deckers JW, Nieboer D, et al. Sex differences in lifetime risk and first manifestation of cardiovascular disease: prospective population based cohort study. *BMJ.* 2014; 349:g5992.
- [26] Santana-Cabrera L, Lorenzo-Torrent R, Sánchez-Palacios M, Martín Santana JD, Hernández Hernández JR. Influence of age in the duration of the stay and mortality of patients who remain in an Intensive Care Unit for a prolonged time. *Rev Clin Esp (Barc).* 2014; 214(2):74-8.
- [27] Gavilanes EG, Gavilanes NL, Fleming R. The effect of age on length of stay in the medical intensive care unit. *Chest.* 2010; 138(4):225A.
- [28] Peigne V, Somme D, Guérot E, Lenain E, Chatellier G, Fagon JY, et al. Treatment intensity, age and outcome in medical ICU patients: results of a French administrative database. *Ann Intensive Care.* 2016; 6(1):7.
- [29] Alisher N, Khokhar RA, Rehman MU, Shaikh AS, Bux H, Sangi R. Impact of cardiopulmonary bypass time and aortic cross clamp time on immediate PostOperative outcomes in patients with congenital heart disease undergoing open heart surgery. *Pak Armed Forces Med J.* 2023; 73(2):443.
- [30] Székely A, Sági E, Király L, Szatmári A, Dinya E. Intraoperative and postoperative risk factors for prolonged mechanical ventilation after pediatric cardiac surgery. *Paediatr Anaesth.* 2006; 16(11):1166-75.
- [31] Remadi JP, Rakotoarivelo Z, Marticho P, Benamar A. Prospective randomized study comparing coronary artery bypass grafting with the new mini-extracorporeal circulation Jostra System or with a standard cardiopulmonary bypass. *Am Heart J.* 2006; 151(1):198.
- [32] Schöttler J, Lutter G, Böning A, Soltau D, Bein B, Caliebe D, et al. Is there really a clinical benefit of using minimized extracorporeal circulation for coronary artery bypass grafting?. *Thorac Cardiovasc Surg.* 2008; 56(02):65-70.
- [33] Polito A, Patorno E, Costello JM, Salvin JW, Emani SM, Rajagopal S, et al. Perioperative factors associated with prolonged mechanical ventilation after complex congenital heart surgery. *Pediatr Crit Care Med.* 2011; 12(3):e122-6.
- [34] Xu S, Liu J, Li L, Wu Z, Li J, Liu Y, et al. Cardiopulmonary bypass time is an independent risk factor for acute kidney injury in emergent thoracic aortic surgery: a retrospective cohort study. *J Cardiothorac Surg.* 2019; 14(1):90.