

Endotracheal Intubation of COVID-19 Patients

Zahid Hussain Khan*

Department of Anesthesiology and Critical Care, Tehran University of Medical Sciences, Tehran, Iran.

The novel coronavirus (Covid-19) emerged for the first time in China and then rapidly spread and swept the entire world like a tornado killing thousands of patients around the planet. People were advised to stay in-doors to prevent the spread of this deadly disease, and this slogan helped to a greater extent in containing the spread of the virus [1].

Unfortunately, there is no treatment for the disease at present but extensive research is going on to find a definitive treatment.

Regarding endotracheal intubation (ETI) of Covid-19 patients, data are scarce and no randomized clinical trials are available to develop and formulate succinct and acceptable guideline in tackling the problem of ETI in these highly risky and vulnerable patients.

Zuo et al. [2] provide us with a step-wise approach in intubating the patients with Novel coronavirus disease. They emphasize that these patients should be intubated employing the videolaryngoscope. In the same vein, Beverley A O [3] does not recommend the use of glidescope for all intubations, but rather suggest that anesthesiologists consider using a glidescope in preference to fiberoptic bronchoscopy in COVID-19 patients.

The use of a glidescope or a similar device is useful but we have some concerns regarding the use of videolaryngoscopes (VLs) based on our own experience from Iran which is probably the worst hit by this deadly disease in the Middle East countries with a high death toll.

Although we agree that VLS should preferentially be used in intubating COVID-19 patients, nevertheless would like to emphasize that ETI with VLs is not always straightforward and easy.

Tracing back the history of ETI ever since the Macintosh direct laryngoscopy (DL) was introduced into clinical practice, it has since that time remained the gold standard for ETI [4]. It requires a direct line of sight to align the three axes (oral-pharyngeal-laryngeal) for optimal glottic visualization, and slight manipulations are needed.

Although new airway devices such as VLS have been introduced, the time to successful intubation had been the shortest when the classic Macintosh laryngoscope was used by anesthesiologists and residents compared to the different forms of VLs [5]. Similarly, others have also floated the idea that it entails a longer time for tracheal intubation when the Glide Scope and McGrath VLs were used in both manikins and patients [6].

Difficulty in passing the endotracheal tube despite improved glottic visualization especially with angulated blade, increased the intubation time [7].

We do agree that videolaryngoscopy is customarily employed when a difficult airway is anticipated. This idea is grounded on the valid assumption that videolaryngoscopy improves the laryngeal view as compared to DL in patients with suspected difficult intubation, but it is shrouded in darkness whether videolaryngoscopy translates to increase in overall endotracheal tube success rates [8].

Furthermore, videolaryngoscopy could possibly be associated with an overall higher rate of complications such as damage of the laryngeal structures including the cuneate and corniculate cartilages as the endotracheal tube is negotiated into the glottic orifice. Such events are noticed when videolaryngoscopy is conducted by a novice or else hurriedly executed by an experienced health provider.

Even with improved glottis visualization, tracheal intubation can still be challenging with varying success rate depending on the provider's experience with the videolaryngoscopy is conducted by a novice or else hurriedly executed by an experienced health provider.

Even with improved glottis visualization, tracheal intubation can still be challenging with varying success rate depending on the provider's experience with the videolaryngoscopy [9]. We have also noticed in our own experience that at times it may be exceedingly difficult to introduce the endotracheal tube through the glottic aperture despite adequate exposure, and the process can only be successfully accomplished if the endotracheal

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: Khanzh51@yahoo.com

Copyright © 2024 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

tube is rotated 90 degrees anticlock wise to avoid its impingement on the cartilages. Moreover, cases of difficult tracheal tube insertion can be managed successfully if smaller size tubes are used or a malleable guide is introduced into the endotracheal tube prior to its insertion through the glottic opening [10].

Zuo et al. [2] emphasis in performing a rapid airway an assessment often neglected in stressful conditions is commendable because such an assessments which hardly takes a few minutes, if not performed can usher in disastrous repercussions if an accidental difficult intubation is encountered. To the airway indexes that they have provided, we would like to add the upper lip bite test [11] which can be easily and quickly executed in the non- edentulous population.

In conclusion, although we fully endorse that a videolaryngoscope with disposable blades should be available in the airway cart, nevertheless feel that it would be prudent and in the best interest of the patients to also ensure the availability of a Macintosh laryngoscope with disposable blades in case the care provider is welltrained with this device in executing the task of ETI under stressful conditions.

References

- [1] Khan ZH, Rahimi M, Makarem J. Stay Indoors: The Slogan for Nowruz 2020. *Arch Neurosci.* 2020; 7(2):e103043.
- [2] Zuo MZ, Huang YG, Ma WH, Xue ZG, Zhang JQ, Gong YH, et al. Chinese Society of Anesthesiology Task Force on Airway Management; Airway Management Chinese Society of Anesthesiology Task Force on. Expert Recommendations for Tracheal Intubation in Critically ill Patients with Novel Coronavirus Disease 2019. *Chin Med Sci J.* 2020; 35(2):105–9.
- [3] Orser BA. Recommendations for Endotracheal Intubation of COVID-19 Patients. *Anesth Analg.* 2020; 130(5):1109-1110.
- [4] Macintosh RR. A new laryngoscope. *Lancet.* 1943; 241:205.
- [5] Pieters BM, Wilbers NE, Huijzer M, Winkens B, van Zundert AA. Comparison of seven videolaryngoscopes with the Macintosh laryngoscope in manikins by experienced and novice personnel. *Anaesthesia.* 2016;71(5):556-64.
- [6] Hodd JA, Doyle DJ, Gupta S, Dalton JE, Cata JP, Brewer EJ, James M, Sessler DI. A mannequin study of intubation with the AP advance and GlideScope Ranger videolaryngoscopes and the Macintosh laryngoscope. *Anesth Analg.* 2011; 113(4):791-800.
- [7] Sun DA, Warriner CB, Parsons DG, Klein R, Umedaly HS, Moulton M. The GlideScope Video Laryngoscope: randomized clinical trial in 200 patients. *Br J Anaesth.* 2005 Mar;94(3):381-4.
- [8] Lee RA, van Zundert AA. Ensuring direct laryngoscopy will not become an extinct skill. *Acta Anaesthesiol Scand.* 2012; 56(6):803.
- [9] Wong E, Ng YY. The difficult airway in the emergency department. *Int J Emerg Med.* 2008; 1(2):107-11.
- [10] Hussain Z, Tavakoli F. Difficult tracheal tube insertion: A new phraseology. *S Afr Med J.* 2016; 106(2):129.
- [11] Khan ZH, Kashfi A, Ebrahimkhani E. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *Anesth Analg.* 2003; 96(2):595-9.