



Can Cerebral Oximetry Near-Infrared Spectroscopy Strategy in Hypothermic Cardiopulmonary Bypass in Congenital Cardiac Surgery be Considered an Advanced Nursing Practice?

Maryam Shakiba¹, Mahmood Hosseinzadeh Maleki², Ghazaleh Khademian³, Amene Ghanbari⁴, Fatemeh Hajipour⁴, Mohamad Amin Younessi Heravi⁵, Saeideh Imani Moghaddam³, Mohsen Yaghubi^{6*}

¹Department of Anesthesiology, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran.

²Department of Cardiac Surgery, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran.

³Department of Critical Care, Razavi Hospital, Mashhad, Iran.

⁴Department of Extra-Corporeal Circulation (ECC), Rajaie Cardiovascular Medical and Research Center, Iran University of Medical Sciences, Tehran, Iran.

⁵Department of Radiology and Medical Physics, North Khorasan University of Medical Sciences, Bojnurd, Iran.

⁶Department of Extra-Corporeal Circulation (ECC), Razavi Hospital, Mashhad, Iran.

ARTICLE INFO

Article history:

Received 18 March 2024

Revised 12 April 2024

Accepted 27 April 2024

Keywords:

Cardiopulmonary bypass;
Cardiac surgery;
Induced hypothermia;
Near infra-red spectroscopy

ABSTRACT

Background: The importance of cerebral perfusion during congenital cardiac surgery, especially after hypothermic cardiopulmonary bypass (CPB) strategy, has remained a debate. In this study, we evaluated the effect of mild hypothermic CPB strategy on cerebral oximetry near-infrared spectroscopy in congenital heart defects surgery.

Methods: In a randomized prospective study, the pediatrics aged two months and six years with RACHS categories I-II for congenital heart defects surgery were randomly divided into normothermic and mild hypothermic CPB groups. The NIRS was measured in all patients via the INVOS™ cerebral oximetry system. The perfusion and anesthesia strategies during the study were followed as standard methods. A p-value below 0.05 is considered to be a significant level.

Results: Regarding cerebral oximetry, the results showed no significant difference between normothermic and mild hypothermic CPB groups.

Conclusion: It can be concluded that optimizing cerebral oxygen saturation monitoring during congenital cardiac surgery can improve patient outcomes as a protective strategy.

Introduction

Cardiopulmonary bypass (CPB) is indispensable to most cardiac surgeries [1]. Many protective strategies have been introduced to decrease CPB

use complications [2]. Nevertheless, none can provide an optimal situation to protect organs from perfusion disturbances [3].

Hypothermic CPB is a protective strategy to diminish the incidence chance of Ischemic- reperfusion injury (I/R

The authors declare no conflicts of interest.

*Corresponding author.

E-mail address: n.m.yaghubi@gmail.com



Injury) [4]. However, the significant consequences of this approach may be neglected [5].

The balance of the cerebral oxygen supply versus oxygen metabolic demand represents cerebral oxygen saturation [6]. Hence, cerebral oximetry Near-infrared spectroscopy (NIRS) is an applied technology for estimating non-invasive systemic oxygen delivery (DO₂) adequacy [7]. Although some evidence demonstrates that this index measures cerebral venous oxyhemoglobin saturation and reflects the adequacy of tissue oxygen delivery during cardiac surgery, the accuracy of this technology in hypothermic situations during cardiac surgery is debated [8].

Methods

Study design and Participants:

This study is a single-center, prospective, randomized cohort study. Patients were randomly allocated to receive either normothermic CPB (35-37°C) or mild hypothermic (28-32°C) CPB strategy.

The patients were all of pediatrics between two months and six years old. They underwent cardiopulmonary bypass (CPB) with risk adjustment for congenital heart surgery (RACHS) categories I-II for congenital heart defects surgery from January 2019 to June 2022, at the Department of cardiac surgery, Imam-Reza Hospital, Mashhad, Iran.

Exclusion criteria were congenital heart surgery (RACHS) categories higher than II, critical conditions (severe renal and hepatic failure) before surgery, upper airway congenital anomalies, the history of Cold Agglutinins, Parents/guardians do not provide written informed consent and cardiac surgeries that require deep hypothermic circulatory arrest (18-28°C). Also, drop-out criteria were intraoperative complications and the need for reoperation.

The sample size was 120 patients (60 patients in the mild hypothermic perfusion and 60 patients in the normothermic group) by considering the means of space and hypothesis of the same study [6], 99% confidence level, 80% power, and weaning time in the mild hypothermic and normothermic groups. Finally, a further 10% was added to anticipate drop-out cases. The method of block randomization, after the patients were deemed eligible and informed consent had been obtained from the parents, was used.

A computer using block randomization of four patients created a randomization schedule. The study managers placed the group assignments in sequentially numbered envelopes and arranged the facilities and data collection.

Perfusion Strategy

All the pediatrics received the same perfusion strategy in both groups. The CPB flow in normothermic group patients was maintained at 3.5 L/min/m² (body surface area), and in mild hypothermic group patients, it was between 2.63 and 3.15 L/min/m².

Non-pulsatile perfusion with a roller pump was recommended. During CPB, the alpha-stat strategy was administered to evaluate arterial blood gas. Thus, PaCO₂ and PaO₂ were maintained at 35-45mmHg and 150-250 mmHg, respectively.

Anesthesia strategy

The anesthetic induction and maintenance were administered in the operation room using a standardized protocol in both groups. Also, mild hypothermia and normothermia were monitored via a nasopharyngeal thermometer.

At first, demographic variables (age, gender, and BMI), cyanotic or acyanotic congenital cardiac defects, and the type of cardiac defects were recorded.

The core endpoint of this research was the near-infrared cerebral oximetry measurements by INVOS™ system (Medtronic, Minneapolis, MN, USA).

Statistical analysis was performed using SPSS software version 26.0 (Chicago, IL, USA). The quantitative results were presented as mean± SD and median (inter-quartile range=IQR) for normal and non-normally distributed data. Categorical data were expressed as frequency (percentage). Normal distribution of the quantitative data was checked using the Shapiro-Wilks test, Q-Q, and Box plots. Independent Student's T-test was used for variables with normal distribution, and the Mann-Whitney test compared non-normal quantitative variables. The homogeneity of categorical variables between the groups was analyzed using the independence Chi-square or Fisher's exact test. After checking relevant assumptions, paired quantitative variables were compared with paired sample t-test or Wilcoxon test. After checking the normality of residuals, in order to investigate the measurement time (repeated measure variable) and interaction of measurement time and categorical group variables, the generalized estimating equation (GEE) method was used. The significance level was considered as P<0.05.

Results

In this study, 120 (50% in the normothermic group and 50% in the mild hypothermic group) eligible patients with a mean± age of 18.6±17.6 months who underwent congenital cardiac defects surgery participated. The male patients comprised 54.2% of all of the patients.

1) Baseline characteristics:

The comparison of the demographic variables of the pediatrics is provided in (Table 1).

The two groups in this study had no significant differences in age, BMI index, type of surgery, and cyanosis or acyanotic status (P>0.05). Only the gender of the patients was heterogeneous between treatment groups (p<0.001).

As the evaluation of the primary outcome, the NIRS evaluation analysis revealed no statistical differences between normothermic CPB and mild hypothermic CPB groups. Nevertheless, the investigation of the effect of the

study groups and measurement time on the NIRS variations through Generalized Estimating Equations (GEE) showed that the normothermic group had a more

significant effect on the time measurement trend than the mild hypothermic CPB group ($p=0.030$).

Table 1- Compares the baseline characteristics between mild hypothermic and normothermic group patients

Variables		Hypothermia group	Normothermia group	P value
Age (mean± SD) Month		19.4±18.3	17.8±16.9	0.629
BMI (mean± SD)		14.3±2.4	14.1±2.7	0.667
Gender; N(%)	Male	43(71.7)	22(36.7)	<0.001*
	Female	17(28.3)	38(63.3)	
Type of Surgery; N(%)	ASD	4(6.7)	6(10)	0.167
	VSD	28(46.7)	36(60)	
	TOF	28(46.7)	18(30)	
Cyanotic disorder; N (%)	Yes	11(18.3)	9(15)	0.624
	No	49(81.7)	51(85)	

*=significant level at $p<0.05$. (ASD= Atrial Septal Defect, VSD= Ventricular Septal Defects, TOF= Tetralogy of Fallot)

Discussion

Currently, systemic hypothermia during CPB among pediatric cardiovascular surgeons is prevalent. Many prefer maintaining core body temperature moderately hypothermic during surgery [9-10]. On the other hand, the normothermic CPB strategy was widely practiced in adult cardiac surgeries, and this perfusion approach was neglected in pediatrics since there was less strong evidence about clinical outcomes in this group [11].

The main finding of this study revealed that the NIRS evaluations had no statistical differences between hypothermic and normothermic CPB strategies. Since this index measures cerebral venous oxyhemoglobin saturation and reflects the adequacy of tissue oxygen delivery, both study groups revealed good tissue perfusion. However, using isolated NIRS for perfusion adequacy is not a sensitive index. It cannot predict other clinical deterioration outcomes, such as neurological disturbance and even low cardiac output states, after pediatric cardiac surgeries [2, 8, 12].

Nevertheless, there is some evidence that this strategy can prevent some post-cardiac surgery neurological complications, especially delirium [13]; the etiology of neurocognitive dysfunction after cardiac surgery with CPB remained a challenging phenomenon [14]. Furthermore, the mechanism of delayed cognitive impairments and recovery in pediatric patients after cardiac surgery is unknown [15].

So, it seems that more clinical investigation with close follow-up of the pediatrics after cardiac surgery in hypothermic and normothermic strategies is crucial.

Conclusion

Although monitoring any physiological parameter can provide vast information to nurses, perfusionists, and some medical groups during and after pediatric cardiac surgery, ensuring cerebral perfusion in pediatrics undergoing hypothermic as in normothermic CPB

strategy may be necessary. Also, providing and applying an optimal strategy for cerebral perfusion monitoring in this group in the perioperative period can potentially impact patient outcomes and enhance their care.

References

- [1] Fathi M, Valaei M, Ghanbari A, Ghasemi R, Yaghubi M. Comparison of Patient's Kidney Function Based on Kidney Disease Improving Global Outcomes (KDIGO) Criteria and Clinical Parameters in Isolated Coronary Artery Bypass Graft (CABG) Surgery in On-Pump and Off-pump Methods in Patients with Low Cardiac Output Syndrome (LCOS) After Surgery. *Anesth Pain Med.* 2020; 10(2):e100517.
- [2] Arki M, Moeinipour A, Fathi M, Abbasi Teshnizi M, Pasandi H, Heydari M. The effects of St. Thomas I and St. Thomas II Cardioplegia solutions on coronary sinus lactate in mitral valve surgery; A Randomized, Double-blinded, Clinical Trial. *Razavi Int J Med.* 2023; 11(3): 12-16.
- [3] Amouzeshi A, Amouzeshi Z, Abbasi Teshnizi M, Moeinipour AA, Hosseinzadeh Maleki M. Off-Pump Versus On-Pump Coronary Artery Bypass Graft Surgery Outcomes During 6 Years: A Prospective Cohort Study. *Acta Med Iran.* 2017; 55(9):578-584.
- [4] Hatami F, Valizadeh N, Salehi F, Hosseinzadeh Maleki M. Topical versus low-dose systemic tranexamic acid in pediatric cardiac surgery: A randomized clinical study. *J Card Surg.* 2020; 35(12):3368-3373.
- [5] Bianco V, Kilic A, Aranda-Michel E, Dunn-Lewis C, Serna-Gallegos D, Chen S, et al. Mild hypothermia versus normothermia in patients undergoing cardiac surgery. *JTCVS Open.* 2021; 7:230-242.
- [6] Ali J, Cody J, Maldonado Y, Ramakrishna H. Near-Infrared Spectroscopy (NIRS) for Cerebral and Tissue Oximetry: Analysis of Evolving Applications. *J Cardiothorac Vasc Anesth.* 2022;

- 36(8 Pt A):2758-2766.
- [7] Loomba RS, Rausa J, Sheikholeslami D, Dyson AE, Farias JS, Villarreal EG, et al. Correlation of Near-Infrared Spectroscopy Oximetry and Corresponding Venous Oxygen Saturations in Children with Congenital Heart Disease. *Pediatr Cardiol.* 2022; 43(1):197-206.
- [8] Rescoe E, Tang X, Perry DA, Sleeper LA, DiNardo JA, Kussman BD, et al. Cerebral near-infrared spectroscopy insensitively detects low cerebral venous oxygen saturations after stage I palliation. *J Thorac Cardiovasc Surg.* 2017; 154(3):1056-1062.
- [9] Xiong T, Pu L, Ma YF, Zhu YL, Cui X, Li H, et al. Safety of Normothermic Cardiopulmonary Bypass in Pediatric Cardiac Surgery: A System Review and Meta-Analysis. *Front Pediatr.* 2021; 9:757551.
- [10] Floh AA, Das S, Haranal M, Laussen PC, Crawford-Lean L, Fan CS, et al. Comparison between Del Nido and conventional blood cardioplegia in pediatric open-heart surgery. *Perfusion.* 2023; 38(2):337-345.
- [11] Haranal M, Chin HC, Sivalingam S, Raja N, Mohammad Shaffie MS, Namasiwayam TK, et al. Safety and Effectiveness of Del Nido Cardioplegia in Comparison to Blood-Based St. Thomas Cardioplegia in Congenital Heart Surgeries: A Prospective Randomized Controlled Study. *World J Pediatr Congenit Heart Surg.* 2020; 11(6):720-726.
- [12] Gagnon MH, Kussman BD, Zhou L, DiNardo JA, Kheir JN. Sensitivity of a Next-Generation NIRS Device to Detect Low Mixed Venous Oxyhemoglobin Saturations in the Single Ventricle Population. *Anesth Analg.* 2020; 131(3):e138-e141.
- [13] Hogue CW, Levine A, Hudson A, Lewis C. Clinical Applications of Near-infrared Spectroscopy Monitoring in Cardiovascular Surgery. *Anesthesiology.* 2021; 134(5):784-791.
- [14] Lozano-Vicario L, Zambom-Ferraresi F, Zambom-Ferraresi F, de la Casa-Marín A, Ollo-Martínez I, Sáez de Asteasu ML, et al. Effectiveness of a multicomponent exercise training program for the management of delirium in hospitalized older adults using near-infrared spectroscopy as a biomarker of brain perfusion: Study protocol for a randomized controlled trial. *Front Aging Neurosci.* 2022; 14:1013631.
- [15] Easley RB, Marino BS, Jennings J, Cassedy AE, Kibler KK, Brady KM, et al. Impaired cerebral autoregulation and elevation in plasma glial fibrillary acidic protein level during cardiopulmonary bypass surgery for CHD. *Cardiol Young.* 2018; 28(1):55-65.